





### NUTRITION

#### Anaemia: Children & Pregnant Women




Source: NFHS, 2005-06

#### Children with birth weight <2.5 KG




Source: AHS

#### Breastfeeding Practices




Source: AHS

#### Equity: Children with birth weight less than 2.5 KG




Source: RSOC, 2013-14

#### Equity: Children (aged 0-5 months) exclusively breastfed




Source: RSOC, 2013-14

#### Stunting, Wasting, Underweight



Source: RSOC, 2013-14

#### Equity: Children under 5 years - Stunted, Wasted, Underweight




Source: RSOC, 2013-14

#### Battling Anaemia amongst mothers and its impacts

Source: AHS, 2012-13

##### Mothers who consumed IFA for 100 days or more (%)




**Odisha: 31.2%**

Highest: Jagatsinghpur (57.5%)

Lowest: Bhadrak (19.5%)

##### Children with birth weight less than 2.5 kgs (%)




**Odisha: 21.6%**

Highest: Koraput (33.0%)

Lowest: Dhenkanal (14.2%)

##### Children (6-35 months) exclusively breastfed for at least 6 months (%)



**Odisha: 30.7%**

Highest: Kandhamal (70.9%)

Lowest: Anugul (15.9%)

Children (6-35 months) exclusively breastfed for at least 6 months (%)